

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 9 hospitals, Jan - Jun 2019 (Region 4)

Organism	TOTAL ISOLATES	BETA - LACTAMS												CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE
<i>Salmonella</i> , Typhoidal			-					-	-	-						-	-							-	-	
<i>Salmonella</i> , Non-Typhoidal	313		45.6 (307)					-	-	81.5 (81)						69.9 (256)	-							-	85.7 (308)	
<i>Shigella</i> spp.	3		-													-	-							-	-	
<i>Staphylococcus aureus</i>		-									-					-	-		-	-	-	-	-	-	-	-
<i>Vibrio cholerae</i> (all serotypes)			-																					-	-	-
<i>Vibrio cholerae</i> O1			-																					-	-	-
<i>Vibrio cholerae</i> non O1, non O139			-																					-	-	-
<i>Vibrio parahaemolyticus</i>	19		-	-	-	-	-	-	-		-					-	-		-					-	-	
<i>Vibrio</i> spp.	3		-	-	-	-	-	-	-		-					-	-		-					-	-	

