

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, RMsC 4, 9 hospitals, Jan - Dec 2017

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES			AMINOGLYCOSIDES		MISCELLANEOUS		
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFZOLIN (U)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	OFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non-typhoidal	911		41.9 (890)											64.9 (188)	100 ^e (47)	. ^e					82.5 (905)	
<i>Vibrio parahaemolyticus</i>	52		 ^e	. ^e		97.9 (48)	94.3 (35)
<i>Vibrio</i> spp.	80		 ^e	. ^e		97.4 (76)	92.9 (56)

