

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 9 hospitals, Jan - Dec 2018 (RMSc4)

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN/ SULBACTAM	PIPERACILLIN/ TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella, Non-Typhoidal</i>	538		(42) (493)					-	-	94.1 (101)						(78) (349)	-							-	(83) (502)		
<i>Shigella spp.</i>			-													-	-							-	-		
<i>Staphylococcus aureus</i>		-									-					-	-		-	-	-	-	-	-	-	-	
<i>Vibrio cholerae (all serotypes)</i>			-																					-	-	-	
<i>Vibrio cholerae non O1, non O139</i>			-																					-	-	-	
<i>Vibrio parahaemolyticus</i>	56		-	-	-	-	-	-	-		-	-	-	-	-	100 (36)	-	-	-						97.2 (36)	96.9 (32)	
<i>Vibrio spp.</i>			-	-	-	-	-	-	-		-	-	-	-	-	-	-	-	-					-	-		

