

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, 9 hospitals, Jan - Dec 2019 (RMSc4 Saraburi)

Organism	TOTAL ISOLATES	BETA - LACTAMS													CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS				
		PENICILLIN	AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Typhoidal			-					-	-	-						-	- <sup>o</sup>							-	-		
<i>Salmonella</i> , Non-Typhoidal	889		43.9 (804)					-	-	89.7 (291)						75.5 (613)	- <sup>o</sup>							-	-		
<i>Shigella</i> spp.	6		-													-	-							-	-		
<i>Staphylococcus aureus</i>		-									- <sup>f</sup>					-	-		-	- <sup>o</sup>	-	-	-	-	-	-	
<i>Vibrio cholerae</i> (all serotypes)	2		-																					-	-	-	
<i>Vibrio cholerae</i> O1			-																					-	-	-	
<i>Vibrio cholerae</i> non O1, non O139	2		-																					-	-	-	
<i>Vibrio parahaemolyticus</i>	61		3.1 (32)	-	-	-	- <sup>o</sup>	-	-	-	-	-	-	-	-	62.5 (40)	-	-	-						-	100 (42)	
<i>Vibrio</i> spp.	16		-	-	-	-	- <sup>o</sup>	-	-	-	-	-	-	-	-	-	-	-	-					-	-		

